

RESEARCH

Open Access



# Increased behavioral problems in children with sleep-disordered breathing

Eszter Csábi<sup>1\*</sup>, Veronika Gaál<sup>2</sup>, Emese Hallgató<sup>1</sup>, Rebeka Anna Schulcz<sup>3</sup>, Gábor Katona<sup>4</sup> and Pálma Benedek<sup>4</sup>

## Abstract

**Background:** Healthy sleep is essential for the cognitive, behavioral and emotional development of children. Therefore, this study aimed to assess the behavioral consequences of sleep disturbances by examining children with sleep-disordered breathing compared with control participants.

**Methods:** Seventy-eight children with SDB (average age: 6.7 years (SD = 1.83); 61 had OSA and 17 had primary snoring) and 156 control subjects (average age: 6.57 years (SD = 1.46) participated in the study. We matched the groups in age ( $t(232) = 0.578, p = 0.564$ ) and gender ( $\chi^2(1) = 2.192, p = 0.139$ ). In the SDB group, the average Apnea-Hypopnea Index was 3.44 event/h (SD = 4.00), the average desaturation level was 87.37% (SD = 6.91). Parent-report rating scales were used to measure the children's daytime behavior including Attention Deficit Hyperactivity Disorder Rating Scale, Strengths and Difficulties Questionnaire, and Child Behavior Checklist.

**Results:** Our results showed that children with SDB exhibited a higher level of inattentiveness and hyperactive behavior. Furthermore, the SDB group demonstrated more internalizing (anxiety, depression, somatic complaints, social problems) ( $p < 0.001$ ) and externalizing (aggressive and rule-breaking behavior) problems compared with children without SDB, irrespective of severity.

**Conclusions:** Based on our findings we supposed that snoring and mild OSA had a risk for developing behavioral and emotional dysfunctions as much as moderate-severe OSA. Therefore, clinical research and practice need to focus more on the accurate assessment and treatment of sleep disturbances in childhood, particularly primary snoring, and mild obstructive sleep apnea.

**Keywords:** Sleep-disordered breathing, Behavior dysregulation, Hyperactivity, Inattentiveness children, Snoring, Sleep apnea

## Background

Sleep-disordered breathing (SDB) is characterized by a broad spectrum of pathology, ranging from partial upper airway obstruction such as primary snoring, to complete upper airway obstruction such as obstructive sleep apnea (OSA). Primary snoring is at the mildest end of the spectrum and is defined as habitual snoring without

respiratory events, abnormalities of gas exchange, or evidence of sleep disruption [1]. OSA is described as prolonged and intermittent upper airway obstruction during sleep resulting in hypoxia, hypercapnia and fragmented sleep [1]. The prevalence of snoring in children ranges from 5 to 12%, of which 5% have OSA [2]. Moreover, Castronovo et al. [3] found that the prevalence of SDB is up to 34.5% [3]. Several medical conditions contribute to the development of SDB such as adenotonsillar hypertrophy, obesity, craniofacial malformation (e.g., Crouzon syndrome), neuromuscular disorders (e.g., muscle hypotonia) or any anatomical abnormalities that narrow the upper airway (e.g., tongue base hypertrophy) [4]. There is

\*Correspondence: eszter.csabi.psy@gmail.com

<sup>1</sup> Department of Cognitive and Neuropsychology, Institute of Psychology, University of Szeged, Szeged, Hungary  
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

emerging evidence that SDB is associated with a deficit in cognitive functioning such as attention [5], information processing [6], executive functions [7–9] and memory [10–12]. These decrements can lead to decreased learning abilities, lower general intelligence [13] and poor academic performance which are frequently observed in children with SDB [14, 15].

Sleep fragmentation and oxygen deprivation might cause cognitive dysfunction by disturbing neuronal myelination and normal blood gas which impacts the development of the brain, particularly in the hippocampus and frontal lobe structures [7, 16, 17]. Previous studies found that the frontal lobe is uniquely vulnerable to the pathological effect of SDB [7, 16]. The frontal lobe develops throughout childhood and is crucial for executive functions which are responsible for higher-order human functioning, self-regulation, inhibition or social interaction. Thus, damage to this region during the maturation period could affect behavioral functioning [18, 19].

The problem with behavioral regulation exhibited by children with SDB also implies frontal lobe dysfunction. Cross-sectional studies have found that children with SDB frequently demonstrate inattention, hyperactivity [20, 21], externalizing problems such as aggression or rule-breaking behavior [6, 22–24] and internalizing problems including anxiety, depression, somatic complaints [5, 21, 24] or social problems [5]. Previous studies suggest that behavioral problems are irrespective of severity [23, 24].

Recent studies supposed that sleepiness might also affect the regulation of emotions and thus could result in impulsivity, irritability and hyperactive behavior [4, 25]. Chervin et al. [25] revealed that snoring and excessive daytime sleepiness related to increased inattention and hyperactive behavior could likely lead to attention-deficit/hyperactivity disorder (ADHD). A meta-analysis by Sedky et al. [20] demonstrated that children with SDB are at a higher risk of presenting ADHD symptoms, such as inattention and hyperactivity. From another perspective, children with ADHD appear to exhibit more sleep difficulties (e.g., delayed sleep onset, bedtime resistance, tiredness during wakefulness) and symptoms of SDB [26].

The current study aims to investigate the behavioral consequences of SDB compared with control participants without SDB. We hypothesized that children with SDB exhibited more behavioral problems than the age-matched control group regardless of the severity of SDB.

## Methods

### Participants

Seventy-eight children with SDB participated in the SDB group. They were recruited from amongst patients who visited the sleep laboratory of the hospitals. All of them

were untreated before the study. Children with a history of developmental, psychiatric or neurological disorders or who were on medication known to affect sleep were excluded. After the physical examination, all of them underwent overnight polygraphy, which was performed using the Somnomedics SOMNOscreen plus device (Randersacker, Germany) at the Sleep Disorders Laboratory of Heim Pal National Pediatric Institute, Budapest, Hungary, and at the Department of Pediatrics, Clinical Center, University of Pécs, Hungary. SDB was diagnosed by a board-certified sleep physician. Patients who met the International Classification of Sleep Disorders criteria [1] for SDB were included in the study. Furthermore, we used the OSA18 Questionnaire to measure the quality of life and to identify the daytime and nighttime symptoms of SDB [27]. The average age of the SDB group was 6.7 years (SD)=1.83) (minimum 4 to maximum 10 years; 32 females/46 males), of which 61 had OSA and 17 had primary snoring. The average Apnea–Hypopnea Index was 3.44 event/h (SD=4.00) (minimum 0 to maximum 19.3 event/h), the average desaturation level was 87.37% (SD=6.91) (minimum 60% to maximum 99%), and the average body mass index was 17.13 kg/m<sup>2</sup> (SD=6.78). According to previous studies, primary snoring has been linked to daytime functional and behavioral impairments [5, 14, 25]. Therefore, we compared the performance of the SDB group to that of the controls and did not examine the OSA and snoring subgroups separately.

The control group consisted of 156 children (average age: 6.57 years (SD=1.46), 80 females/76 males). The control group was recruited through collaboration with public schools. We matched the two groups on age ( $t(232)=0.578$ ,  $p=0.564$ ) and gender ( $\chi^2(1)=2.192$ ,  $p=0.139$ ). They did not suffer from any developmental, psychiatric, or neurological disorders and were free of any sleeping disorders.

Based on the OSA18 questionnaire [27] we excluded those children from the control group who reported SDB-like symptoms and reached at least 60 points or more on the questionnaire. The comparison of the SDB and control groups is presented in Table 1.

### Behavioral assessment

#### *Attention Deficit Hyperactivity Disorder Rating Scale (ADHD-RS)*

ADHD-RS is an 18-item rating scale designed to assess symptoms of inattention, impulsivity and hyperactivity disorder. Each item is rated on a 4-point scale. The rating scale consists of two subscales based on the relevant symptoms: Inattention problems and Hyperactivity-Impulsivity [28]. We used the parent rating scale version in the study.

**Table 1** Mean Ranks and medians of OSA-18 subscales

	SDB group <sup>a</sup>		Control group <sup>b</sup>		P
	Mean Ranks	Median	Mean Ranks	Median	
Sleep Disorders	195.5	15	78,5	4	<.001
Physical Distress	175.56	15	88.47	7	<.001
Diurnal Problems	163.31	8,5	94.59	4	<.001
Emotional Distress	165.86	10	93.32	4	<.001
Caretaker Preoccupation	181.47	13,5	85.51	4	<.001
Total score of OSA-18	192.8	72	79.85	34	<.001

<sup>a</sup> N = 78, <sup>b</sup> N = 156

### **Strengths and Difficulties Questionnaire (SDQ)**

SDQ is a 25-item rating scale used to measure emotional and behavioral problems. Each item is rated on a 3-point scale. The SDQ has five subscales, four of them measure major difficulties commonly experienced by children (Hyperactivity, Emotional Symptoms, Behavioral Problems, Social Problems). One subscale assesses strengths (Prosocial Behavior). There are three versions of the scale: parent-report, teacher-report and self-report versions. This study was based on the parent-report version of the scale [29].

### **Child Behavior Checklist (CBCL)**

CBCL is designed to assess behavioral and emotional problems and competencies. The Hungarian version of the questionnaire consists of 46 items, each item is rated on a 3-point scale. The CBCL measures two factors: Internalizing, composed of subscales Anxious/Depressed, Somatic Complaints, Inattention Problems and Social Problems; and Externalizing, consisting of Aggressive Behavior, Deviant/Rule-Breaking Behavior. The CBCL is available in three versions: parent report, teacher report, and self-report versions. The parent version of the scale was used [30].

### **OSA18 questionnaire**

OSA-18 is a caregiver-administrated quality of life survey that contains 18 items [27]. The questionnaire is organized into five domains: sleep disorders, physical distress, emotional distress, diurnal problems and caretaker preoccupation. Each domain can be scored on a 7-point Likert scale. The OSA-18 total score is the sum of all 18 items. According to Franco et al. [27], a total score < 60, between 60 and 80, and above 80 suggests a small impact on the quality of life, moderate impact and large impact, respectively.

### **Polysomnographic measurement**

The study was performed using the Somnomedics SOMNOscreen plus device (Somnomedics) according to the guideline. Polygraphic readings were evaluated by a physician experienced in sleep medicine. Breathing irregularities (apneas and hypopneas) were analyzed and Apnea–Hypopnea Index (AHI) was presented as the number of apneas and hypopneas per sleeping hour. The desaturation index indicates the number of periods with desaturation (minimum of 3% fall in oxygen saturation) per hour. According to the polysomnographic diagnostic criteria for childhood OSA by Marcus et al. [1] one episode of apnea during sleep is considered pathologic. The respiratory pause has to last for at least two breaths to be considered abnormal. In hypopnea, the respiratory amplitude is reduced by at least 30% and occurs when the oxygen saturation drops by 3%. We classified the severity of OSAS as mild, AHI = 1 event/hour with desaturation or AHI 2–5 event/hour; moderate, 5 < AHI > 10 event/hour; and severe AHI > 10 event/hour [31].

### **Procedure**

All children with SDB underwent a standard pediatric overnight clinical polygraphy. Medical histories and physical examinations were conducted by a board-certified sleep physician before the diagnostic night. On the night of the study, the parents, most often the mothers, completed the demographic and behavioral questionnaires. Written informed consent was obtained from both parents, and procedures were verbally explained to the children before the commencement of the study. The participants did not receive any financial compensation for their participation. Ethics approval was obtained by the Ethics Committee at Heim Pal National Pediatric Institute, Budapest (number of the Ethical Approval: KUT-6/2017).

**Statistical analysis**

Statistical analysis was performed using the SPSS 22.0 (Statistical Package for the Social Sciences). Gender proportions were compared between the groups using Pearson’s chi-square test. Group differences were tested with the Mann–Whitney U test. Spearman’s correlation was used to explore the relationship between respiratory variables and behavioral measurements.

**Results**

The figure represents medians: a) ADHD-RS Rating Scale: Attentional Problem (AP), Hyperactivity-Impulsivity (HI). b) Strength and Difficulty Questionnaire (SDQ): Hyperactivity (H), Emotional Problems (ES), Behaviour Problem (BP), Social Problem (SP), Prosocial Behaviour (PB). c) Total Score of SDQ (TSS). d) Child Behavior Checklist (CBCL): Social Problems (SP), Anxious/Depressed (AD), Rule-Breaking Behavior (RBB), Aggressive Behaviour (AB), Somatic Complaints (SC), Attentional Problems (AP). e) Total Score of CBCL.  $p < 0.001^{***}, p < 0.05^{**}$ .

**ADHD rating scale**

The SDB group demonstrated significantly higher ratings than the control group on both subscales of ADHD-RS: Attentional problems ( $U = 3833, z = -4.638, p < 0.001$ ) and Hyperactivity-Impulsivity ( $U = 3899, z = -4.498, p < 0.001$ ). The medians are presented in Fig. 1.

**Strength and Difficulty Questionnaire**

We revealed a significant effect of group on all SDQ subscales. The SDB group demonstrated significantly higher rates on Hyperactivity ( $U = 4631, z = -2.943, p = 0.003$ ), Emotional Symptoms ( $U = 4548, z = -3.234, p = 0.001$ ),


Behavior problems ( $U = 4437, z = -3.234, p = 0.001$ ) and Social problems ( $U = 4716, z = -2.917, p = 0.004$ ). Furthermore, children with SDB showed significantly lower rates on the subscales of prosocial behavior than the control participants ( $U = 4589, z = -3.136, p = 0.002$ ). The Total number of SDQ was significantly higher in the SDB group than in the control group ( $U = 3852, z = -4.526, p < 0.001$ ) (for medians see Fig. 1).

**Child Behavior Checklist**

The subscales of externalizing ( $U = 4041.500, z = -4.241, p < 0.001$ ) and internalizing problems ( $U = 3893, z = -4.522, p < 0.001$ ) significantly differed between the two groups. The SDB group demonstrated significantly higher scores than the control group on the subscales of Social Problems ( $U = 3954.500, z = -4.449, p < 0.001$ ), Anxious/Depressed ( $U = 4496, z = -3.373, p = 0.001$ ), Aggressive Behavior ( $U = 4149, z = -4.046, p < 0.001$ ) and Rule-Breaking Behavior ( $U = 4480, z = -3.681, p < 0.001$ ). Furthermore, the SDB group had significantly higher rates on Somatic Complaints ( $U = 4524, z = -3.723, p < 0.001$ ) and Attentional Problems ( $U = 3978.500, z = -4.443, p < 0.001$ ). The Total Problems Summary Scale was higher in the SDB group compared to the control group ( $U = 3516, z = -5.269, p < 0.001$ ). The medians are presented in Fig. 1.

**Discussion**

Our goal was to examine the behavioral consequences of SDB compared to control participants without SDB. Our results revealed that children with all severities of SDB demonstrated a significantly higher level of inattention, hyperactivity-impulsivity, internalizing (social problems, anxiety, depression, somatic complaints) and



**Fig. 1** Comparison of the behavioural assessment between SDB and healthy control (HC) groups

externalizing problems (rule-breaking behavior, aggression) than control subjects. Consistent with these results, children with SDB exhibited significantly less prosocial behavior.

Our results are in line with earlier studies showing similar irregular behavioral performance in children with SDB compared to controls, [20–22] irrespective of severity [23, 24, 32]. A study by Rosen et al. [24] also found that increased behavioral morbidity was similar between children with primary snoring and children with OSA. Both groups showed significantly more behavioral problems than the control subjects, such as hyperactivity, emotional lability, somatic complaints, oppositional, aggressive behavior and social problems. The authors revealed that externalizing acting out behavior such as hyperactive, oppositional and aggressive behavior showed the most robust association with SDB [24]. In contrast, some studies supposed that the duration of the disease and severity may contribute to the development of more serious behavioral outcomes [22, 33]. Mulvaney et al. [22] provided evidence that an increased frequency of aggressive behavior, inattention and social problems are related to high Respiratory Disturbance Index (RDI—occurrence of apneas and hypopneas).

Contrary to these findings, Jackman et al. [34] revealed that primary snoring and mild OSA are associated with a higher level of internalizing problems than moderate or severe types of OSA. The authors supposed that the parents of the children diagnosed with mild SDB had more unrelated psychological distress than the parents of children who showed more severe SDB symptoms, which engendered more behavioral problems in the children themselves or caused parents to describe their children's behavior in a more negative light [27]. In line with these results, Lewin et al. [6] demonstrated that children with mild OSA had more symptoms of anxiety, depression, social and externalizing problems than children with severe OSA. The authors suggest that their results might be explained by the fact that the severe OSA group did not show abnormalities in the organization of sleep stages; however, they had significantly more arousals than the mild OSA group [6].

A recent study by Smith et al. [35] suggests that frequent snoring is a more effective predictor of behavioral outcomes than AHI. The authors found that behavioral problems increased with the frequency of snoring up to occasional snoring (at least two nights per week) but did not increase based on AHI severity. These results emphasized the importance of examining snoring in the assessment of potential behavioral outcomes in SDB. To summarize, there has been inconsistency across studies in the relationship between the severity of SDB and behavioral consequences. Therefore, future studies need

to clarify the effect of the severity of SDB on behavioral functioning and the role of individual vulnerability.

The mechanisms causing the behavioral consequences have not been fully delineated. Some investigators suggest that hypoxia impacts the development of the central nervous system, particularly the hippocampus and frontal lobe structures [7, 16]. Others postulated that sleep fragmentation produces behavioral problems [14]. A few studies assumed that daytime sleepiness is associated with impaired behavior, especially hyperactivity and externalizing acting-out behavior like irritability and impulsivity [4, 25]. However, excessive daytime sleepiness is a less common symptom in children with SDB. In children, sleepiness is frequently acted out behaviorally rather than expressed verbally (e. g. complaining of being tired) [4]. Our findings are consistent with those studies that suggest an overlap between the impairment associated with SDB and the diagnostic criteria for ADHD [20]. However, is still unclear whether sleep disturbances are intrinsic to ADHD or sleep disorders cause ADHD-like symptoms. In the future, we need to clarify whether SDB is intrinsic to ADHD or SDB causes ADHD-like symptoms to avoid misdiagnosis. Therefore, children should be examined for SDB when considering a diagnosis of ADHD.

The main limitation of the study is the lack of overnight polygraphic measurement of the control group. With these data, we would have a deeper insight into the differences and effects of pathological and non-pathological ventilation during sleep on daytime functioning. Nonetheless, our results suggest that mild SDB (e. g., snoring) has a risk of developing behavioral and emotional dysfunctions as much as moderate or severe SDB. Overall, this study highlights that clinical practice need to focus more the adequate treatment of sleep disturbances in childhood, particularly primary snoring, and mild obstructive sleep apnea.

## Conclusion

Our findings revealed that children with SDB demonstrated higher levels of emotional (e. g., anxiety, depression, somatic complaints) and behavioral difficulties (e. g., inattentiveness, hyperactive, aggressive and rule-breaking behavior) compared with children without SDB, irrespective of severity. We believe that our study provides evidence that the potential impact of mild SDB may be greater than what was previously believed. Our results confirm the findings of previous studies suggesting that even snoring and a mild level of SDB can increase the risk of behavioral and emotional problems. Our study underlines the importance of early and accurate assessment and adequate treatment of primary snoring as much as moderate or severe type of SDB.

## Abbreviations

SD: Standard Deviation; SDB: Sleep-Disordered Breathing; OSA: Obstructive Sleep Apnea; ADHD: Attention Deficit and Hyperactivity Disorder; ADHD-RS: Attention Deficit Hyperactivity Disorder Rating Scale; SDQ: Strengths and Difficulties Questionnaire; CBCL: Child Behavior Checklist; AHI: Apnea-Hypopnea Index; RDI: Respiratory Disturbance Index.

## Acknowledgements

We are thankful to the subjects for their participation and the nurses of the Sleep Disorders Laboratory of Heim Pal National Pediatric Institute for their work and assistance.

## Authors' contributions

ECS and PB designed the study. The material preparation, data collection was performed by ECS, PB, VG, RS and GK. The statistical analysis was conducted by HE and ECS. The first draft of the manuscript was written by ECS and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

## Funding

Open access funding provided by University of Szeged. The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

## Availability of data and materials

The full data set and other materials related to about this study can be obtained from the corresponding author on reasonable request.

## Declarations

### Ethics approval and consent to participate

The study protocol was reviewed and approved by the ethical review board of Heim Pal National Pediatric Institute, Budapest, Hungary (number of the Ethical Approval: EL4.1-F3) in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments.

**Informed consent:** Informed consent was obtained from all individual participants included in the study.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no conflict of interest.

### Author details

<sup>1</sup>Department of Cognitive and Neuropsychology, Institute of Psychology, University of Szeged, Szeged, Hungary. <sup>2</sup>Department of Pediatrics, Clinical Center, University of Pécs, Pécs, Hungary. <sup>3</sup>Sleep Disorders Laboratory of Heim Pal National Pediatric Institute, Budapest, Hungary. <sup>4</sup>Department of Oto-Rhino-Laryngology and Head-Neck Surgery, University of Szeged, Szeged, Hungary.

Received: 22 November 2021 Accepted: 31 August 2022

Published online: 15 September 2022

## References

- Marcus CL, Marcus Brooks LJ, Davidson Ward S, Draper KA, Gozal D, Halbower AC, Jones J, Lehmann C, Schechter MS, Sheldon S, Shiffman RN, Spruyt K. Diagnosis and Management of Childhood Obstructive Sleep Apnea Syndrome. *Pediatrics*. 2012;130(3):e714–55. <https://doi.org/10.1542/peds.2012-1672>.
- Hangström K, Saarenpää-Heikkilä O, Himanen SL, Lapinlampi AM, Nieminen P. Behavioral problems and neurocognitive functioning in snoring school-aged children. *Psychol Cogn Sci*. 2015;1(2):46–53. <https://doi.org/10.17140/PCSOJ-1-107>.
- Castronovo V, Zucconi M, Nosselt L, Marazzini C, Hensley M, Veglia F, Nespoli L, Ferini-Strambi L. Prevalence of habitual snoring and sleep-disordered breathing in preschool-aged children in an Italian community. *J of Pediatr*. 2003;142:377–82. <https://doi.org/10.1067/mpd.2003.118>.
- Trosman I, Trosman SJ. Cognitive and Behavioral Consequences of Sleep Disordered Breathing in children. *Med Sci*. 2017;5:2–14. <https://doi.org/10.1007/s11325-010-0329-4>.
- Blunden S, Lushington K, Kennedy D, Martin J, Dawson D. Behavior and neurocognitive performance in children aged 5–10 years who snore compared to controls. *J Clin and Exp Neuropsychol*. 2000;22:554–68. [https://doi.org/10.1076/1380-3395\(200010\)22:5;1-9;FT554](https://doi.org/10.1076/1380-3395(200010)22:5;1-9;FT554).
- Lewin DS, Rosen RC, England SJ, Dahl RE. Preliminary evidence of behavioral and cognitive sequelae of obstructive sleep apnea in children. *Sleep Med*. 2002;3:5–13. [https://doi.org/10.1016/S1389-9457\(01\)00070-3](https://doi.org/10.1016/S1389-9457(01)00070-3).
- Beebe DW, Gozal D. Obstructive sleep apnea and the prefrontal cortex: toward a comprehensive model linking nocturnal upper airway obstruction to daytime cognitive and behavioral deficit. *J Sleep Res*. 2002;11:1–16. <https://doi.org/10.1046/j.1365-2869.2002.00289.x>.
- Kohler M, Lushington K, van den Heuvel C, Martin J, Pamula Y, Kennedy D. Adenotonsillectomy and neurocognitive deficits in children with sleep disordered breathing. *PLoS ONE*. 2009;4:e7343. <https://doi.org/10.1371/journal.pone.0007343>.
- Karpinski AC, Scullin MH, Montgomery-Downs HE. Risk for sleep-disordered breathing and executive function in preschoolers. *Sleep Med*. 2008;9(4):418–24. <https://doi.org/10.1016/j.sleep.2007.06.004>.
- Csábi E, Benedek P, Janacsek K, Katona G, Németh D. Sleep-disorder in childhood impairs declarative but not nondeclarative forms of learning. *Clin Exp Neuropsychol*. 2013;35:677–85. <https://doi.org/10.1080/13803395.2013.815693>.
- Csábi E, Benedek P, Janacsek K, Zavec ZS, Katona G, Németh D. Declarative and non-declarative memory consolidation in children with sleep disorders. *Front Hum Neurosci*. 2016;9:709. <https://doi.org/10.3389/fnhum.2015.00709>.
- Kheirandish-Gozal L, DeJong MR, Spruyt K, Chamuleau SAJ, Gozal D. Obstructive sleep apnoea is associated with impaired pictorial memory task acquisition and retention in children. *Eur Respir J*. 2010;36:164–9. <https://doi.org/10.1183/09031936.00114209>.
- Kennedy JD, Blunden S, Hirte C, Parsons DW, Martin AJ, Crowe E, Fracs W, Pamula Y, Lushington K. Reduced neurocognition in children who snore. *Pediatr Pulmonol*. 2004;37:330–7. <https://doi.org/10.1002/ppul.10453>.
- Urschitz MS, Guenther A, Eggebrecht E, Wolff J, Urschitz-Duprat PM, Schlaud M, Poets CF. Habitual snoring, intermittent hypoxia and academic performance in primary school children. *American J Respir Crit Care Med*. 2004;168:464–8. <https://doi.org/10.1164/rccm.200212-1397OC>.
- Chervin RD, Clarke DF, Huffman JL, Szymanska E, Ruzickaa DL, Miller V, Nettlesd AL, Sowerse MFR, Giordan BJ. School performance, race and other correlates of sleep-disordered breathing in children. *Sleep Med*. 2003;4:21–7. [https://doi.org/10.1016/s1389-9457\(02\)00243-5](https://doi.org/10.1016/s1389-9457(02)00243-5).
- Halbower AC, Degaonkar M, Barker PB, Earley CJ, Marcus CL, Smith L, Prahme MC, Mahone EM. Childhood obstructive sleep apnea associates with neuropsychological deficits and neuronal brain injury. *Plos Med*. 2006;3: e301. <https://doi.org/10.1371/journal.pmed.0030301>.
- Chan KL, Shi L, So HK, Wang D, Liew AWC, Rasalkar DD, Chu CW, Wing JK, Li AM. Neurocognitive dysfunction and grey matter density deficit in children with obstructive sleep apnea. *Sleep Med*. 2014;15:1055–61. <https://doi.org/10.1016/j.sleep.2014.04.011>.
- Landau YE, Bar-Yishay O, Greenberg-Dotan S, Goldbart AD, Tarasiuk A, Asher T. Impaired behavioral and neurocognitive functioning preschool children with obstructive sleep apnea. *Pediatr Pulmol*. 2012;47:180–8. <https://doi.org/10.1002/ppul.21534>.
- O'Brien LM. The neurocognitive effects of sleep disruption in children and adolescents. *Sleep Med Clin*. 2011;6:109–16. <https://doi.org/10.1016/j.jsmc.2010.12.007>.
- Sedky K, Bennett DS, Carvalho KS. Attention deficit hyperactivity disorder and sleep disorder breathing in pediatric populations: a meta-analysis. *Sleep Med Rev*. 2014;18:349–56. <https://doi.org/10.1016/j.jsmrv.2013.12.003>.
- Ting H, Hong Wong R, Yang HJ, Lee SP, Lee SD, Wang L. Sleep-disordered breathing, behavior and academic performance in Taiwan schoolchildren. *Sleep Breath*. 2011;15:91–8. <https://doi.org/10.1007/s11325-010-0329-4>.
- Mulvaney SA, Goodwin JL, Morgan WJ, Rosen GR, Quan SF, Kaemingk KL. Behavior problems associated with sleep disordered breathing in

- school-aged children – the Tucson Children's Assessment of Sleep Apnea Study. *J Ped Psychol*. 2005;31:322–30. <https://doi.org/10.1093/jpepsy/jsj035>.
23. Bourke RS, Anderson V, Yang JSC, Jackman AR, Killedar A, Nixon GM, Davey MJ, Walker AM, Trinder J, Horne RSC. Neurobehavioral function is impaired in children with all severities of sleep disordered breathing. *Sleep Med*. 2011;12:222–9. <https://doi.org/10.1016/j.sleep.2010.08.011>.
  24. Rosen CL, Storfer-Isser A, Taylor HG, Kirchner HL, Emancipator JL, Redline S. Increased behavioral morbidity in school-aged children with sleep-disordered breathing. *Pediatrics*. 2004;114:1640–8. <https://doi.org/10.1542/peds.2004-0103>.
  25. Chervin RD, Archbold KH, Dillon JE, Panahi P, Pituch KJ, Dahl RE, Guilleminault C. Inattention, hyperactivity and symptoms of Sleep-Disordered Breathing. *Pediatr*. 2002;109(3):449–56. <https://doi.org/10.1542/peds.109.3.449>.
  26. Weiss MD, Salpekar J. Sleep problems in child with Attention-Deficit Hyperactivity Disorder. *CNS Drugs*. 2010;24(10):1–18. <https://doi.org/10.2165/11538990-000000000-00000>.
  27. Franco AR, Rosenfeld RM, Rao M. First place—resident clinical science award 1999. Quality of life for children with obstructive sleep apnea. *Otolaryngol Head Neck Surg*. 2000;123:9–16. <https://doi.org/10.1067/mhn.2000.105254>.
  28. DuPaul GJ, Anastopoulos AD, Power TJ, Reid R, Ikeda MJ, McGoey KE. Parent Ratings of Attention-Deficit/Hyperactivity Disorder Symptoms: Factor Structure and Normative Data. *J Psychopathol Behav Assess*. 1998;20:83–102. <https://doi.org/10.1023/A:1023087410712>.
  29. Goodman R. Psychometric Properties of the Strengths and Difficulties Questionnaire. *J Am Acad Child Adolescent Psychiatry*. 2001;40(11):1337–45. <https://doi.org/10.1097/00004583-200111000-00015>.
  30. Achenbach TM, Howell CT, Quay HCC, Conners CK, Bates JE. National Survey of Problems and Competencies among Four- to Sixteen-Year-Olds: Parents' Reports for Normative and Clinical Samples. *Monogr Soc Res Child Dev*. 1991;56(3):1–135. <https://doi.org/10.2307/1166156>.
  31. Berry RB, Budhiraja R, Gottlieb DJ, Gozal D, Iber C, Kapur VK, Marcus CL, Mehra R, Parthasarathy S, Quan SF, Redline S, Strohl KP, Davidson Ward SL, Tangredi MM. Rules for scoring respiratory events in sleep: update of the 2007 AASM manual for the scoring of sleep and associated events. Deliberations of the sleep apnea definitions task force of the American academy of sleep medicine. *J of Clin Sleep Med*. 2012;8(5):597–619. <https://doi.org/10.5664/jcsm.2172>.
  32. Brockmann PE, Urschitz MS, Schlaud M, et al. Primary snoring in school children: prevalence and neurocognitive impairments. *Sleep Breath*. 2012;16(1):23e9. <https://doi.org/10.1007/s11325-011-0480-6>.
  33. Ersu R, Arman AR, Save D, Karadag B, Karakoc F, Berkem M, Dagli E. Prevalence of snoring and symptoms of sleep-disordered breathing in primary school children in istanbul. *Chest*. 2004;126:19–24. <https://doi.org/10.1378/chest.126.1.19>.
  34. Jackman AR, Biggs SN, Walter LM, Embuldeniya US, Davey MJ, Nixon GM, Anderson V, Trinder J, Horne RSC. Sleep-disordered breathing in preschool children is associated with behavioral but not cognitive impairments. *Sleep Med*. 2012;13:621–31. <https://doi.org/10.1016/j.sleep.2012.01.013>.
  35. Smith DL, Gozal D, Hunter J, Kheirandish-Gozal L. Frequency of snoring, rather than apnea hypopnea index, predicts both cognitive and behavioral problems in young children. *Sleep Med*. 2017;34:170–8.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

